
Opioid Overdose Medical History Form (derived from 20 ILCS 301/5-23)

Name of OD Responder _____ Date _____

The following questions are to be answered about the person the naloxone is intended for

___ Myself ___ Friend / partner ___ Family member: _____ ___ Other: _____

Name _____ Age _____ M F Race/Ethnicity _____

Age first used: Any opiates _____ Daily opiates _____ Needle _____

Currently use >1-2 times/month: Cocaine? _____ Alcohol? _____ Speed? _____ Benzos? _____

How many times in: Detox (<7d) _____ Rehab (>7d) _____ Jail _____ Methadone _____

Number of times: Self Detoxed _____ Self OD'd _____ Witness OD _____ Witness OD death _____

Taking any other medications? _____

NALOXONE EDUCATION CHECKLIST & STANDING ORDER TO DISTRIBUTE – OD Responder should initial when complete

_____ **OD prevention:** Educate OD prevention partners; know your source; purity testing; tie-release “tasting”

_____ **Risk Factors:** mixing drugs (including cocaine!), period of abstinence for any reason (as little as 3 days)

_____ --S-- **Signs of Overdose:** unresponsive to arousal / pain; slower breathing; blue lips.

_____ --C-- **Call 911!!**

_____ --A-- **A & B of Life:** clear **Airway** of food / gum / etc. **Breath** rate >6-7/min. Recovery position

_____ --R-- **Rescue Breathing:** Position; clear airway; pinch off nose; 2 big quick breaths then breathe every 5 seconds ... but meantime have someone preparing naloxone!

_____ --E-- **Evaluate...** Is it a good time to inject naloxone?

_____ --M-- **Naloxone/narcan:** KEEP IT HANDY!! Draw 1cc into syringe w/1 in+ needle; inject into muscle! If nasal naloxone is being used, put device together as instructed and inject into nostrils. If Evzio is being used, follow all of the voice prompts. Continue rescue breathing until the person wakes up. If they aren't up in 1-2 minutes, give another dose.

_____ --E-- **Aftercare:** Naloxone/narcan works for 30-40 min. Support them from using again! Naloxone/narcan wears off in 30-60 min, so OD on long-acting opioids (methadone, oxy, etc.) may return
-- be ready to give second dose and/or transport to hospital

OD Educator

Date

Prescriber
